

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/04448 A2

(51) International Patent Classification⁷: C07D 473/00

(21) International Application Number: PCT/US01/21373

(22) International Filing Date: 6 July 2001 (06.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/216,844 7 July 2000 (07.07.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
NEOTHERAPEUTICS, INC. [US/US]; 157 Technology Drive, Irvine, CA 92618 (US).

Published:

— without international search report and to be republished upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DIAMOND, Jack** [CA/CA]; 25 Chedoke Avenue Lane, Hamilton, Ontario L8P 4B1 (CA). **GLASKY, Alvin, J.** [US/US]; 11955 Lambert, Tustin, CA 92782 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(74) Agents: **CULLMAN, Louis, C.** et al.; Oppenheimer Wolff & Donnelly LLP, Suite 700, 840 Newport Center Drive, Newport Beach, CA 92660-7007 (US).



WO 02/04448 A2

(54) Title: METHODS FOR TREATMENT OF DRUG-INDUCED PERIPHERAL NEUROPATHY AND RELATED CONDITIONS

(57) Abstract: A method of treating drug-induced peripheral neuropathy comprises administering to a patient with drug-induced peripheral neuropathy an effective quantity of a purine derivative or analogue, a tetrahydroindolone derivative or analogue, or a pyrimidine derivative or analogue. If the compound is a purine derivative, the purine moiety can be guanine or hypoxanthine. The compound can induce peripheral nerve sprouting through the action of a neurotrophic factor such as nerve growth factor (NGF) without the occurrence of hyperalgesia. The peripheral nerve sprouting can be nociceptive nerve sprouting. The drug-induced peripheral neuropathy can be drug-induced peripheral neuropathy associated with the administration of oncolytic drugs, such as a vinca alkaloid, cisplatin, paclitaxel, suramin, altretamine, carboplatin, chlorambucil, cytarabine, dacarbazine, docetaxel, etoposide, fludarabine, ifosfamide with mesna, tamoxifen, teniposide, or thioguanine. Methods according to the present invention are particularly useful in treating peripheral neuropathy associated with the administration of vincristine, paclitaxel, or cisplatin.

**METHODS FOR TREATMENT OF DRUG-INDUCED
PERIPHERAL NEUROPATHY AND RELATED CONDITIONS**

CROSS-REFERENCES

This application claims priority from Provisional Application Serial No.

- 5 60/216,844, filed July 7, 2000 by Jack Diamond and Alvin J. Glasky, and entitled "Methods for Treatment of Peripheral Neuropathy and Related Conditions with Bifunctional Purine Analogues," which is incorporated herein in its entirety by this reference.

BACKGROUND OF THE INVENTION

- 10 This invention is directed to methods for treatment of drug-induced peripheral neuropathy and related conditions, particularly drug-induced peripheral neuropathy associated with the administration of oncolytic drugs.

- Many oncolytic or antineoplastic drugs have been developed in recent years. Although such drugs have proven effective in many cases in the treatment of
- 15 malignancies, they can have severe side effects. One of the most serious and clinically significant side effect is peripheral neuropathy. Many antineoplastic drugs can cause peripheral neuropathy. For some of the most effective drugs, neurotoxicity is dose-limiting. It can force the termination of otherwise successful therapy, or can preclude the repetition of successful therapy. Sensory abnormalities produced by the
- 20 administration of antineoplastic drugs can range from mild paresthesiae or dysesthesiae to severe neuropathic pain. In some cases, sensory and motor symptoms resolve within days or weeks after the agents are discontinued. However, peripheral neuropathy can be a chronic painful and disabling condition. The mechanisms that produce peripheral neuropathy as a consequence of the
- 25 administration of oncolytic drugs are largely unknown (R.C. Polomano & G.J. Bennett, "Chemotherapy-evoked Painful Peripheral Neuropathy," Pain Med. 2: 8-14 (2001); S. De Santis et al., "Patients Treated with Antitumor Drugs Displaying Neurological Deficits Are Characterized by a Low Circulating Level of Nerve Growth Factor," Clin. Cancer Res. 6: 90-95 (2000); K. Hayakawa et al., "NGF Prevention of Neurotoxicity
- 30 Induced by Cisplatin, Vincristine and Taxol Depends on Toxicity of Each Drug and NGF Treatment Schedule: In Vitro Study of Adult Rat Sympathetic Ganglion Explants," Brain Res. 794: 313-319 (1998)).

Accordingly, there is a need for more efficient methods of combating drug-induced peripheral neuropathy, particularly peripheral neuropathy induced by the administration of oncolytic drugs. Preferably, such methods should not interfere with cancer treatment or block the activity of the oncolytic drugs. Such methods should also not induce other side effects and should be well tolerated by cancer patients. Preferably, such methods should also combat peripheral neuropathy for all oncolytic drugs and should not depend on specific interactions with each individual oncolytic drug. There is a particular need for methods that can stimulate nerve growth or regeneration, particularly without inducing hyperalgesia.

10 SUMMARY

One embodiment of the present invention is a method of treating drug-induced peripheral neuropathy comprising administering to a patient with drug-induced peripheral neuropathy an effective quantity of an effective quantity of a compound comprising: (1) a moiety A selected from the group consisting of a purine moiety, a purine analogue, a tetrahydroindolone moiety, a tetrahydroindolone analogue, a pyrimidine moiety, a pyrimidine analogue and aminocarboxaminoimidazoles; (2) a hydrocarbyl moiety L of 1 to 6 carbon atoms that is linked to the moiety A and that can be cyclic, with the hydrocarbyl moiety being optionally substituted with one or more substituents selected from the group consisting of lower alkyl, amino, hydroxy, lower alkoxy, lower alkylamino, lower alkylthio, and oxo; and (3) a moiety B that is linked to the moiety L through a carbonyl group wherein B is -OZ or N(Y₁)-D, where Z is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, aralkyl, or heteroaralkyl; D is a moiety that promotes absorption of the compound; and Y₁ is hydrogen, alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms, which can be N, O, or S.

The purine moiety can be selected from the group consisting of hypoxanthine and guanine, as well as other purine moieties. A number of purine derivatives suitable for use in methods according to the present invention are disclosed. In addition to

these purines, analogues of naturally occurring purines defined above, tetrahydroindolones, tetrahydroindolone analogues, pyrimidines, pyrimidine analogues and aminocarboxaminoimidazoles, which maintain the proposed pharmacophore between the 6, 5, 4, and 9 positions on the purine molecule, may be used as alternatives to purines.

Preferably, the compound is capable of passing through the blood-brain barrier.

Typically, the administration of the compound induces peripheral nerve sprouting in the skin of the patient to whom the purine derivative is administered. The peripheral nerve sprouting can be nociceptive nerve sprouting. Typically, the nociceptive nerve sprouting is induced without the occurrence of hyperalgesia.

The drug-induced peripheral neuropathy can be induced by the administration of an oncolytic drug such as a vinca alkaloid, cisplatin, paclitaxel, suramin, altretamine, carboplatin, chlorambucil, cytarabine, dacarbazine, docetaxel, etoposide, fludarabine, ifosfamide with mesna, tamoxifen, teniposide, or thioguanine. Methods according to the present invention are particularly significant in treating drug-induced peripheral neuropathy arising from the administration of vincristine, paclitaxel, or cisplatin.

BRIEF DESCRIPTION OF THE DRAWINGS

The following invention will become better understood with reference to the specification, appended claims, and accompanying drawings, where:

Figure 1 is a bar graph showing the effects of riluzole and AIT-082 in the prevention of cisplatin toxicity in DRG cells;

Figure 2 is a graph showing the effects of vincristine or vincristine plus AIT-082 on body weight;

Figure 3 is a bar graph showing sensory nerve conduction velocity demonstrating the effects of AIT-082 on vincristine-induced neuropathy;

Figure 4 is a bar graph showing EMG amplitude demonstrating the effects of AIT-082 on vincristine-induced neuropathy;

Figure 5 is a bar graph showing the effect of AIT-082 on the total number of myelinated nerve fibers after vincristine administration.

DESCRIPTION

We have discovered that the bifunctional purine derivative N-4-carboxyphenyl-3-(6-oxohydropurin-9-yl) propanamide (also known as AIT-082 and leteprinin potassium), which bypasses the blood-brain barrier, can act to induce peripheral nerve sprouting in the skin of adult rats. As detailed below in the Example, this activity may be attributable to upregulation of cutaneous nerve growth factor (NGF) levels induced by this bifunctional purine derivative, although Applicants do not intend to be bound by this theory. Moreover, this activity occurred without the induction of hyperalgesia. This property of acting to induce peripheral nerve sprouting, therefore, should also be possessed by other purine derivatives and analogues, tetrahydroindolone derivatives and analogues, and pyrimidine derivatives and analogues, as discussed below.

The peripheral nerve sprouting can be nociceptive nerve sprouting. The nociceptive nerve sprouting can occur without the induction of hyperalgesia.

Typically, a compound useful in a method of the present invention is capable of bypassing the blood-brain barrier.

More specifically, as detailed below in the Example, systematically administered AIT-082 closely mimics the effects both of increased levels of endogenous NGF, and of exogenous NGF. The compound induces vigorous collateral sprouting but has no effect on axonal regeneration after nerve crush (shown earlier to occur independently of NGF), and the sprouting it induced was blocked by systemic anti-NGF treatment. The growth of such nerve tissue is evoked and maintained entirely by the increased levels of NGF in adjacent denervated skin. However, AIT-082 resembles more the effects of increased endogenous NGF than of exogenous NGF, because it did not induce hyperalgesia. In fact, the sprouting evoked by AIT-082 was shown to be entirely attributable to the measured up-regulation it induced in endogenous NGF levels in skin.

Accordingly, one aspect of the present invention is a method of treating drug-induced peripheral neuropathy comprising administering to a patient with drug-induced peripheral neuropathy an effective quantity of a compound, the compound comprising:

- (1) a moiety A selected from the group consisting of a purine moiety, a purine analogue, a tetrahydroindolone moiety, a tetrahydroindolone analogue, a pyrimidine moiety, and a pyrimidine analogue; (2) a hydrocarbyl moiety L of 1 to 6 carbon atoms that is linked to the moiety A and that can be cyclic, with the hydrocarbyl moiety being optionally substituted with one or more substituents selected from the group consisting


of lower alkyl, amino, hydroxy, lower alkoxy, lower alkylamino, lower alkylthio, and oxo; and (3) a moiety B that is linked to the moiety L through a carbonyl group wherein B is $-OZ$ or $N(Y_1)-D$, where Z is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, aralkyl, or heteroaralkyl; D is a moiety that promotes absorption of the compound having activity
 5 against a multi-drug transporter protein; and Y_1 is hydrogen, alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl,
 10 heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms, which can be N, O, or S.

Typically, a compound useful in a method of the present invention is capable of passing through the blood-brain barrier.

15 In one preferred embodiment of methods according to the present invention, the moiety A is a purine moiety.

In one alternative, A is a substituted or unsubstituted hypoxanthine moiety. Typically, in this alternative, L has the structure $-(CH_2)_n$ where n is an integer from 1 to 6.


20 The compound having the activity against drug-induced peripheral neuropathy can be a compound of formula (I)



where n is an integer from 1 to 6 and R is hydrogen or lower alkyl or is a salt or prodrug ester of a compound of formula (I) wherein n is an integer from 1 to 6 and R is
 25 hydrogen or lower alkyl. Typically, the compound is a compound of formula (I) wherein n is an integer from 1 to 6 and R is hydrogen or lower alkyl. Typically, R is hydrogen, and the compound is N-4-[[3-(6-oxo-1,6-dihydropurin-9-yl)-1-oxopropyl]

amino] benzoic acid, designated AIT-082. Alternatively, R is ethyl, and the compound is N-4-[[3-(6-oxo-1,6-dihydropurin-9-yl)-1-oxopropyl] amino] benzoic acid ethyl ester.

When the purine moiety is hypoxanthine, a preferred purine derivative is a compound of formula (I)




5

(I)

wherein n is an integer from 1 to 6 or of a salt or prodrug ester of formula (I) wherein n is an integer from 1 to 6. Typically, the purine derivative is a compound of formula (I) wherein n is an integer from 1 to 6. Preferably, n is 2 and the compound is N-4-carboxyphenyl-3-(6-oxohydropurin-9-yl) propanamide, also known as AIT-082. The activity of this compound is described further in the Example.

Alternatively, the purine derivative can be a 9-substituted hypoxanthine derivative of formula (II)




(II)

wherein n is a integer from 1 to 6, R₁ is selected from the group consisting of H, COOH, and COOW₁, where W₁ is selected from the group consisting of lower alkyl, amino, and lower alkylamino, and R₂ is selected from the group consisting of H and OH.

In this alternative, for one particularly preferred purine derivative, n is 2, R₁ is H and R₂ is OH and the purine derivative is N-(2-(5-hydroxyindol-3-yl))ethyl-3-(6-

oxohydropurine-9-yl) propanamide. In this alternative, for another particularly preferred purine derivative, n is 2, R₁ is H and R₂ is H and the purine derivative is N-(2-indol-3-yl)ethyl-3-(6-oxohydropurin-9-yl) propanamide. In this alternative, for still another particularly preferred purine derivative, n is 2, R₁ is COOH, and R₂ is OH and the purine derivative is N-(1-carboxyl-2-(5-hydroxyindol-3-yl))ethyl-3-(6-oxohydropurin-9-yl) propanamide.


As another alternative, the purine derivative can be a 9-substituted hypoxanthine derivative of formula (III)



wherein n is an integer from 1 to 6, R₁ is selected from the group consisting of H, COOH, and COOW₁, wherein W₁ is selected from the group consisting of lower alkyl, amino, and lower alkylamino, R₂ is selected from the group consisting of H and OH, and R₃ is selected from the group consisting of H and OH.

In this alternative, for one particularly preferred purine derivative, n is 2, R₁ is H, R₂ is H, and R₃ is OH, and the purine derivative is N-(2-(3,4-dihydroxyphenyl))ethyl-3-(6-oxohydropurin-9-yl) propanamide. In this alternative, for another particularly preferred purine derivative, n is 2, R₁ is H, R₂ is OH, and R₃ is OH, and the purine derivative is N-(2-hydroxy-2-(3,4-dihydroxyphenyl))ethyl-3-(6-oxohydropurin-9-yl) propanamide. In this alternative, for still another particularly preferred purine derivative, n is 2, R₁ is COOH, R₂ is H, and R₃ is OH, and the purine derivative is N-(1-carboxyl-2-(3,4-dihydroxyphenyl))ethyl-3-(6-oxohydropurin-9-yl) propanamide.

When the purine moiety is guanine, one preferred purine derivative is a 9-substituted guanine derivative of formula (IV)




(IV)

wherein n is an integer from 1 to 6, R_1 is selected from the group consisting of H, COOH, and COOW₁, or W₁ is lower alkyl, amino, or lower alkylamino, and R_2 is selected from the group consisting of H and OH.

- 5 In this alternative, for one particularly preferred purine derivative, n is 2, R_1 is H, and R_2 is OH, and the purine derivative is N-(2-(5-hydroxyindol-3-yl))ethyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide. In this alternative, for another particularly preferred purine derivative, n is 2, R_1 is H, and R_2 is H and the purine derivative is N-(2-(2-indol-3-yl)ethyl))-3-(2-amino-6-oxohydropurin-9-yl)) propanamide. In this alternative, for still
- 10 another particularly preferred purine derivative, n is 2, R_1 is COOH, and R_2 is OH, and the purine derivative is N-(1-carboxyl)-(2-(5-hydroxyindol-3-yl))ethyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide.


Alternatively, the purine derivative can be a 9-substituted guanine derivative of formula (V) wherein n is an integer from 1 to 6.



(V)

15 In this alternative, for one particularly preferred purine derivative, n is 2 and the compound is N-4-carboxyphenyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide.


Alternatively, the purine derivative can be a 9-substituted guanine derivative of formula (VI) wherein n is an integer from 1 to 6.



(VI)

In this alternative, for one particularly preferred purine derivative, n is 2 and the compound is 3-(2-amino-6-oxohydropurine-9-yl) propanoic acid.


Alternatively, the purine derivative can be a 9-substituted guanine derivative of formula (VII) wherein n is an integer from 1 to 6, p is an integer from 1 to 6, and q is an integer from 1 to 3.



(VII)

In this alternative, for one particularly preferred purine derivative, n is 2, p is 2, and q is 1, and the purine derivative is N-[2-[[2-(2-oxopyrrolidin-1-yl)-1-oxoethyl]amino]ethyl] propanamide.

Alternatively, the purine derivative can be a 9-substituted guanine derivative of formula (VIII) wherein R_1 is selected from the group consisting of H, COOH, and COOW₁, where W₁ is selected from the group consisting of lower alkyl, amino, and lower alkylamino, R_2 is selected from the group consisting of H and OH, and R_3 is selected from the group consisting of H and OH.




(VIII)

In this alternative, for one particularly preferred purine derivative, n is 2, R_1 is H, R_2 is H, and R_3 is OH, and the purine derivative is N-(2-(3,4-dihydroxyphenyl)ethyl)-3-

(2-amino-6-oxohydropurin-9-yl) propanamide. In this alternative, for another particularly preferred purine derivative, n is 2, R₁ is H, R₂ is OH, and R₃ is OH, and the purine derivative is N-(2-hydroxy-2-(3,4-dihydroxyphenyl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide. In this alternative, for still another particularly preferred purine derivative, n is 2, R₁ is COOH, R₂ is H, and R₃ is H and the compound is N-(1-carboxyl-2-(3,4-dihydroxyphenyl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.


Alternatively, the purine derivative can be a 9-substituted guanine derivative of formula (IX) wherein n is an integer from 1 to 6 and p is an integer from 1 to 3.



In this alternative, for one particularly preferred purine derivative, n is 2, p is 1, and the compound is the 1-(dimethylamino)-2-propyl ester of N-4-carboxyphenyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

Other bifunctional hypoxanthine derivatives suitable for use in methods according to the present invention are disclosed in U.S. Patent No. 5,091,432 to Glasky, incorporated herein by this reference. Other bifunctional guanine derivatives suitable for use in methods according to the present invention are disclosed in U.S. Patent Application No. 09/49,153, by Glasky et al., incorporated herein by this reference.

More generally, purine-based compounds suitable for use in methods according to the present invention are compounds in which A is a substituted or unsubstituted 9-atom bicyclic moiety in which the 5-membered ring has 1 to 3 nitrogen atoms, the bicyclic moiety having the structure of formula (X)



where:

(1) if the bond between N₁ and the bond between C₅ is a single bond, then the bond between C₆ and R₆ is a double bond, R₆ is O or S, and R₁ is hydrogen, alkyl, aralkyl, cycloalkyl, or heteroaralkyl;

(2) if the bond between N₁ and C₆ is a double bond, then the bond between C₆ and R₆ is a single bond, R₁ is not present, and R₆ is hydrogen, halo, amino, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁, where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(3) if the bond between C₂ and N₃ is a single bond, then the bond between C₂ and R₂ is a double bond, R₂ is O or S, and R₃ is hydrogen or alkyl;

(4) if the bond between C₂ and N₃ is a double bond, then the bond between C₂ is a single bond, R₃ is not present, and R₂ is hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, amino, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁, where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl,

aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, aralkoxy carbonyl, heteroaralkoxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(5) ~~A₇ and A₈ are C or N;~~

(a) if A₇ and A₈ are both C and the bond between A₇ and A₈ is a single bond, then the bond between A₈ and R₈ is two single bonds to two hydrogen atoms or is a double bond in which R₈ is O or S and R₇ is two hydrogen atoms;

(b) if A₇ and A₈ are both C and the bond between A₇ and A₈ is a double bond, then R₇ is hydrogen, the bond between A₈ and R₈ is a single bond and R₈ is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

(c) if A₇ and A₈ are both N, then the bond between A₇ and A₈ is a double bond, and R₇ and R₈ are not present;


(d) if A₇ is C and A₈ is N, then the bond between A₇ and A₈ is a double bond, R₇ is hydrogen, and R₈ is not present;

(e) if A₇ is N, A₈ is C, and the bond between A₇ and A₈ is a double bond, then R₇ is not present, the bond between A₈ is a single bond, and R₈ is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

(f) if A₇ is N, A₈ is C, and the bond between A₇ and A₈ is a single bond, then R₇ is hydrogen, alkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl, the bond between A₈ and R₈ is a double bond, and R₈ is O or S; and

(6) N₉ is bonded to L; with the proviso that A does not have the structure of an unsubstituted guanine or hypoxanthine.

The purine moiety can be a purine moiety of formula (XI)



5 in which:

(1) R₁ is selected from the group consisting of hydrogen, alkyl, aralkyl, cycloalkyl, and heteroaralkyl; and

(2) R₂ is selected from the group consisting of hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁,
 10 where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are
 15 present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, arylkoxycarbonyl, heteroarylokoxycarbonyl,
 20 alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroarylkylaminocarbonyl in which the alkyl portions could be cyclic and can contain from one to three heteroatoms which could be N, O, or S, with the proviso that both R₁ and R₂ are not hydrogen and that R₁ is not hydrogen when R₂ is amino.

25 The purine moiety of formula (XI) is a hypoxanthine or a guanine derivative but excludes unsubstituted hypoxanthine, in which R₁ and R₂ are hydrogen, and unsubstituted guanine, in which R₁ is hydrogen and R₂ is amino.


In one particularly preferred embodiment, R₁ is butyl and R₂ is hydrogen.

In another preferred embodiment, R₁ is benzyl and R₂ is hydrogen.

30 In another preferred embodiment, R₁ is dimethylaminoethyl and R₂ is hydrogen.

In another preferred embodiment, R₁ is cyclopentyl and R₂ is hydrogen.

In another preferred embodiment, R₁ is cyclohexylmethyl and R₂ is hydrogen.
 In another preferred embodiment, R₁ is cyclopropylmethyl and R₂ is hydrogen.
 In another preferred embodiment, R₁ is hydrogen and R₂ is phenyl.
 In another preferred embodiment, R₁ is hydrogen and R₂ is trifluoromethyl.
 In another preferred embodiment, R₁ is hydrogen and R₂ is butyl.
 In another preferred embodiment, R₁ is butyl and R₂ is butyl.
 In another preferred embodiment, R₁ is hydrogen and R₂ is methyl.
 In another preferred embodiment, R₁ is hydrogen and R₂ is phenylamino.
 Alternatively, the purine moiety is a purine moiety of Formula (XII)



in which:

- (1) R₂ is selected from the group consisting of hydrogen, halo, amino, OQ₃, SQ₃, NHNH₂, NHOQ₃, NQ₃Q₄, or NHQ₃, where Q₃ and Q₄ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, and heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₃ and Q₄ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₃ where Y₃ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, aralkoxy carbonyl, heteroaryloxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and
- (2) R₆ is selected from the group consisting of hydrogen, halo, amino, OQ₅, SQ₅, NHNH₂, NHOQ₅, NQ₅Q₆, or NHQ₆, where Q₅ and Q₆ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, and heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1


to 3 heteroatoms which can be N, O, or S, and when Q₅ and Q₆ are present together and are alkyl, they can be taken together to form a 5- or 6- membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, arylkoxycarbonyl, heteroarylkoxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S.

In one preferred example of this embodiment, R₂ is hydrogen and R₆ is -NH₂ or -N(CH₃)₂.

In another preferred example of this embodiment, R₂ is hydrogen and R₆ is Cl.

In yet another preferred example of this embodiment, R₂ is -NH₂ and R₆ is Cl.

In another alternative, the purine moiety is the purine moiety of Formula (XIII)



in which:

(1) R₁ is hydrogen, alkyl, aralkyl, cycloalkyl, or heteroaralkyl; and


(2) R₂ is O or S.

Preferably, in this embodiment, R₁ is hydrogen and R₂ is O or S.

Particularly preferred purine-based compounds for use in methods according to the present invention include: (1) 4-[3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester; (2) 4-[3-(1-butyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester; (3) 4-[3-(1-methyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester; (4) 4-[3-(1-(2-dimethylaminoethyl)-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester; (5) 4-[3-(2,6-dioxo-1,2,3,6-tetrahydropurin-9-yl)propionylamino] benzoic acid ethyl ester; (6) 4-[3-(6-methoxypurin-9-yl)propionylamino] benzoic acid ethyl ester; (7) 4-[3-(6-dimethylaminopurin-9-yl)propionylamino] benzoic acid ethyl ester; (8) 4-[3-(2-amino-6-chloropurin-9-yl)propionylamino] benzoic acid ethyl ester; (9) 4-[2-(6-oxo-2-thioxo-

- 1,2,3,6-tetrahydropurin-9-yl)propionylamino]benzoic acid ethyl ester; (10) 4-[2-(2-butyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino]benzoic acid ethyl ester; (11) 4-[2-(6-oxo-2-phenyl-1,6-dihydropurin-9-yl)propionylamino]benzoic acid ethyl ester; (12) 4-[[3-(6-chloropurin-9-yl)propionyl]methylamino] benzoic acid methyl ester; (13) 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-[3-(2-oxopyrrolidin-1-yl)propyl] propionamide; (14) 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-{2-[2-(2-oxopyrrolidin-1-yl)acetyl]amino}ethyl} propionamide; (15) N-3-(2-oxopyrrolidin-1-yl)propyl]-3-(6-oxo-2-thioxo-1,2,3,6-tetrahydropurin-9-yl) propionamide; and (16) 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-(3-morpholin-4-yl-propyl) propionamide.

- 10 In another alternative of methods according to the present invention, the compound is a tetrahydroindolone derivative or analogue where A is a 9-atom bicyclic moiety in which the 5-membered ring has one to three nitrogen atoms, the bicyclic moiety having the structure of formula (XIV)



- 15 where:

(1) N₁ is bonded to L;

(2) A₂ and A₃ are C or N;

- (a) If A₂ and A₃ are both C and the bond between A₂ and A₃ is a single bond, then the bond between A₂ and R₂ is two single bonds, two hydrogen atoms or is a double bond in which R₂ is O or S and R₃ is two hydrogen atoms;

(b) If A₂ and A₃ are both C and the bond between A₂ and A₃ is a double bond, then R₃ is hydrogen, the bond between A₂ and R₂ is a single bond and R₂ is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

- (c) If A₂ and A₃ are both N, then the bond between A₂ and A₃ is a double bond and R₂ and R₃ are not present;

(d) If A₂ is N and A₃ is C, then the bond between A₂ and A₃ is a double bond, R₂ is not present, and R₃ is hydrogen;

(e) If A_2 is C, A_3 is N, and the bond between A_2 and A_3 is a double bond, then R_3 is not present, the bond between A_2 and R_2 is a single bond, and R_2 is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

5 (f) If A_2 is C, A_3 is N, and the bond between A_2 and A_3 is a single bond, then R_3 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, or heteroaralkenyl, the bond between A_2 and R_2 is a double bond, and A_2 is O or S;

(3) R_5 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NHQ_1 , NQ_1Q_2 , OH, 10 OQ_1 , or SQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, 15 aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_1 and Q_2 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other 20 heteroatom, which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, 25 arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;


(4) R_5 is hydrogen unless R_5 is alkyl, in which case R_5 is hydrogen or the same alkyl as R_5 ;

30 (5) R_5 and R_5 can be taken together as a double bond to C_5 , and can be O, S, NQ_3 , or C which can be substituted with one or two groups R_5 , where Q_3 is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

- (6) R_6 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, NH_2 , NH_4 , NQ_4Q_5 , OH, OQ_4 , or SQ_4 , where Q_4 and Q_5 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_4 and Q_5 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom, which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylamino carbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

- (7) R_6' is hydrogen unless R_6 is alkyl, in which case R_6' is hydrogen or the same alkyl as R_6 ;
- (8) R_6 and R_6' can be taken together as a double bond to C_6 and can be O, S, NQ_6 , or C which can be substituted with one or two groups R_5 , and where Q_6 is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and
- (9) R_7 is hydrogen unless R_5 is alkyl and R_5' is hydrogen, in which case R_7 is the same alkyl as R_5 .

Typically, A is a tetrahydroindolone moiety. More typically, the tetrahydroindolone moiety is a tetrahydroindolone moiety of formula (XV)




in which:

- (1) R_5 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NH_1 , NQ_1Q_2 , OH , OQ_1 , or SQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S;
- (2) R_5 is hydrogen;
- (3) R_6 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NHW_1 , NQ_1Q_2 , OH , OQ_1 , or SQ_1 , where Q_1 and Q_2 are aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S and where W_1 is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S;
- (4) R_6 is hydrogen; and
- (5) R_7 is hydrogen.

Typically, R_5 , R_5' , R_6 , R_6' , and R_7 are all hydrogen.

- When A is a tetrahydroindolone moiety, preferred compounds are 4-[3-(4-oxo-4,5,6,7-tetrahydroindolon-1-yl) propionylamino] benzoic acid ethyl ester and 4-[3-(4-oxo-4,5,6,7-tetrahydroindolon-1-yl) propionylamino] benzoic acid.

In another alternative, the compound is a pyrimidine derivative or pyrimidine analogue. In this alternative, A is an amino-substituted 6-membered heterocyclic moiety of formula (XVI)



where:

- (1) if the bond between N_1 and the bond between C_6 is a single bond, then the bond between C_6 and R_6 is a double bond, R_6 is O or S, and R_1 is hydrogen,

alkyl, aralkyl, cycloalkyl, or heteroaralkyl;

(2) if the bond between N_1 and C_6 is a double bond, then the bond between C_6 and R_6 is a single bond, R_1 is not present, and R_6 is hydrogen, halo, amino, OH, OQ_1 , SQ_1 , $NHNH_2$, NQ_1Q_2 , or NHQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_1 and Q_2 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl,

aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(3) if the bond between C_2 and N_3 is a single bond, then the bond between C_2 and R_2 is a double bond, R_2 is O or S, and R_3 is hydrogen or alkyl;

(4) if the bond between C_2 and N_3 is a double bond, then the bond between C_2 and R_2 is a single bond, R_3 is not present, and R_2 is hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, amino, OH, OQ_1 , SQ_1 , $NHNH_2$, NHQ_1 , NQ_1Q_2 , or NHQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_1 and Q_2 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y_3 , where Y_3 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in

which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(5) R_4 is hydrogen, alkyl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, or heteroarylaminocarbonyl;

(6) A_5 is carbon or nitrogen;

(7) if A_5 is nitrogen, then R_5 is not present;

(8) if A_5 is carbon, then R_5 is hydrogen, amino, alkyl, alkoxy, halo, nitro, aryl, cyano, alkenyl, or alkaryl;

(9) if R_5 and R_6 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and

(10) N_4 is bonded to L.

Typically, A_5 is carbon and the 6-membered heterocyclic moiety is a pyrimidine moiety.

When A is a pyrimidine moiety, in one alternative, R_2 is O and R_3 is hydrogen. In this alternative, the pyrimidine moiety can be cytosine, thymine, uracil, 3-methyluracil, 3-methylthymine, 4-methylcytosine, 5-methylcytosine, 5-hydroxymethylcytosine, 5-hydroxyuracil, 5-carboxymethyluracil, or 5-hydroxymethyluracil.

In another alternative, R_2 is S and R_3 is hydrogen. In this alternative, the pyrimidine moiety can be 2-thiouracil, 5-methylamino-2-thiouracil, 5-methyl-2-thiouracil, or 2-thiocytosine.

In still another alternative, R_2 is amino and the bond between C_2 and N_3 is a double bond. In this alternative, the pyrimidine moiety can be 2-aminopyrimidinone or 2-amino-4-chloropyrimidine.

In still another alternative, R₂ is hydrogen and the bond between C₂ and N₃ is a double bond. In this alternative, the pyrimidine moiety can be 4-chloropyrimidine, 5-amino-4-chloropyrimidine, 4-chloro-5-methylpyrimidine, 4-chloro-5-hydroxymethylpyrimidine, or 4-chloro-5-carboxymethylpyrimidine.

- 5 In still another alternative, R₁ is hydrogen, methyl, or ethyl, R₅ is hydrogen, methyl, or ethyl, and R₆ is O. In this alternative, the pyrimidine moiety can be pyrimidinone.

- Particularly preferred pyrimidine compounds include: 4-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 4-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 4-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 4-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid; 4-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid; 4-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid; 3-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 3-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 3-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester; 3-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid; 3-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid; and 3-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.
- 10
15
20

- In accordance with the present invention, and as used herein, the following terms, when appearing alone or as part of a moiety including other atoms or groups, are defined with the following meanings, unless explicitly stated otherwise. In addition, all groups described herein can be optionally substituted unless such substitution is excluded. The term "alkyl," as used herein at all occurrences, refers to saturated aliphatic groups including straight-chain, branched-chain, and cyclic groups, all of which can be optionally substituted. Preferred alkyl groups contain 1 to 10 carbon atoms. Suitable alkyl groups include methyl, ethyl, and the like, and can be optionally substituted. The term "alkenyl," as used herein at all occurrences, refers to unsaturated groups which contain at least one carbon-carbon double bond and includes straight-chain, branched-chain, and cyclic groups, all of which can be optionally substituted. Preferable alkenyl groups have 2 to 10 carbon atoms. The term "alkoxy" refers to the ether —O—alkyl, where alkyl is defined as as above. The term "aryl" refers to aromatic groups which have at least one ring having a conjugated
- 25
30

π -electron system and includes carbocyclic aryl and biaryl, both of which may be optionally substituted. Preferred aryl groups have 6 to 10 carbon atoms. The term "aralkyl" refers to an alkyl group substituted with an aryl group. Suitable aralkyl groups include benzyl and the like; these groups can be optionally substituted. The term "aralkenyl" refers to an alkenyl group substituted with an aryl group. The term "heteroaryl" refers to carbon-containing 5-14 membered cyclic unsaturated radicals containing one, two, three, or four O, N, or S heteroatoms and having 6, 10, or 14 π -electrons delocalized in one or more rings, e.g., pyridine, oxazole, indole, thiazole, isoxazole, pyrazole, pyrrole, each of which can be optionally substituted as discussed above. The term "sulfonyl" refers to the group $-S(O_2)-$. The term "alkanoyl" refers to the group $-C(O)R_g$, where R_g is alkyl. The term "aroyl" refers to the group $-C(O)R_g$, where R_g is aryl. Similar compound radicals involving a carbonyl group and other groups are defined by analogy. The term "aminocarbonyl" refers to the group $-NHC(O)-$. The term "oxycarbonyl" refers to the group $-OC(O)-$. The term "heteroaralkyl" refers to an alkyl group substituted with a heteroaryl group. Similarly, the term "heteroaralkenyl" refers to an alkenyl group substituted with a heteroaryl group. As used herein, the term "lower," in reference to an alkyl or the alkyl portion of an another group including alkyl, is defined as a group containing one to six carbon atoms. The term "optionally substituted" refers to one or more substituents that can be lower alkyl, aryl, amino, hydroxy, lower alkoxy, aryloxy, lower alkylamino, arylamino, lower alkylthio, arylthio, or oxo, in some cases, other groups can be included, such as cyano, acetoxy, or halo. The term "halo" refers generally to fluoro, chloro, bromo, or iodo; more typically, "halo" refers to chloro.


As indicated above, the linker L is a hydrocarbyl moiety of 1 to 6 carbon atoms that can be cyclic, with the hydrocarbyl moiety being optionally substituted with one or more substituents selected from the group consisting of lower alkyl, amino, hydroxy, lower alkoxy, lower alkylamino, lower alkylthio, and oxo. Preferably, the linker L has the structure $-(CH_2)_n-$ wherein n is an integer from 1 to 6. As detailed below, for most preferred embodiments of compounds useful in methods according to the present invention, a preferred linker has n equal to 2 or 3.

The moiety B is either: (i) $-OZ$, where Z is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, aralkyl, or heteroaralkyl; or (ii) $N(Y_1)-D$, where D is a moiety that promotes absorption of the compound, and Y_1 is hydrogen, alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, which, when taken with D, can form a cyclic 5- or 6-membered

saturated ring which can contain one other heteroatom which can be O, N, or S, of which N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, arylloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S. Typically, Y₁ is hydrogen. Where the moiety B is -OZ, the moiety B is a carboxylic acid or carboxylic acid or ester. Typically, where B is a carboxylic acid ester, the moiety Z is a lower alkyl, such as methyl, ethyl, butyl, propyl, or isopropyl.


In one alternative, the moiety D, as described above, is a moiety having at least one polar, charged, or hydrogen-bond-forming group to improve the metabolic and bioavailability properties of the compound. The moiety D can be, but is not limited to, a moiety with physiological or biological activity such as nootropic activity. In one alternative, the moiety D can be a moiety containing at least one carboxyl, carboxamide, carboxyl ester, or carbonyl function. In another alternative, the moiety D can be a moiety containing at least one hydroxyl, primary amino, secondary amino, tertiary amino, sulfhydryl, or sulfonamidyl function. The moiety D can be cyclic or acyclic. Preferred examples of the moiety D are described below.

When the moiety D is a cyclic or acyclic moiety containing at least one carbonyl, carboxamide, carboxyl ester, or carbonyl function, in one preferred example, D is a carboxylic acid or carboxylic acid ester with the structure




wherein p is an integer from 1 to 6 and W₁ is selected from the group consisting of hydrogen and lower alkyl. Typically, if W₁ is lower alkyl, it is methyl, ethyl, propyl, butyl, or isobutyl. Typically, p is 3. Typically, W₁ is hydrogen or ethyl.

In another preferred example, D and Y₁ are taken together to form a piperazine derivative as described in D. Manetti et al., "Molecular Simplification of 1,4-Diazabicyclo[4.3.0]nonan-9-ones Gives Piperazine Derivatives That Maintain High Nootropic Activity," J. Med. Chem. 43: 4499-4507 ("Manetti et al. (2000)"). B is an analogue of structure




wherein Q₁ is hydrogen, methyl, ethyl, butyl, or propyl, Q₂ is hydrogen or methyl, where, if Q₂ is methyl, it can be located at either of the two possible positions in the piperazine ring.

5 In another preferred example, D has the structure




where one of Z₁ and Z₂ is hydrogen, and the other of Z₁ and Z₂ is -COOH or -COOW₁, wherein W₁ is alkyl. Typically, W₁ is selected from the group consisting of methyl, ethyl, propyl, butyl, and isobutyl. Either of Z₁ or Z₂ can be hydrogen. When Z₁ is hydrogen and Z₂ is -COOH, the moiety B is *p*-aminobenzoic acid (PABA). When Z₁ is -COOH and Z₂ is hydrogen, the moiety B is *m*-aminobenzoic acid (MABA). When Z₁ is hydrogen and Z₂ is -COOW₁, the moiety B is an ester of *p*-aminobenzoic acid (PABA). When Z₁ is -COOW₁ and Z₂ is hydrogen, the moiety B is an ester of *m*-aminobenzoic acid (MABA). Typically, these esters are ethyl esters.

15 When the moiety D is a moiety that contains at least one hydroxyl, primary amino, secondary amino, tertiary amino, sulfhydryl, or sulfonamidyl function, in one preferred example, D is a phenylsulfonamidyl moiety of structure




wherein p is an integer from 0 to 6. Typically, p is 2.

20 In another preferred example, D is an alkylpyridyl moiety of structure



wherein p is an integer from 1 to 6. Typically, p is 1.


In another preferred example, D is a dialkylaminoalkyl moiety of the structure



wherein p is an integer from 1 to 6 and Q₇ and Q₈ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken
 5 together to form a 5 or 6 member ring which may contain 1 other heteroatom which can be N, O, or S, of which the N may be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl,
 10 aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S.

Where Q₇ and Q₈ can be taken together to form a five or six member ring, the
 15 ring is typically pyrrolidine, piperidine, or morpholine. The pyrrolidine ring can be optionally substituted with oxo. The piperidine ring can be optionally substituted with methyl or ethyl. Typically, p is 2 or 3.

In another preferred example, D is an alkylpyrrolidine moiety of the structure



20 wherein p is an integer from 1 to 6 and W₁ is selected from the group consisting of methyl, ethyl, and propyl. Typically, W₁ is methyl. Typically, p is 2.

Preferably, a compound useful in methods according to the present invention has a log P of from about 1 to about 4 in order to optimize bioavailability and CNS penetration of the compound.

25 As detailed below in the Example, compounds used in methods according to the present invention are believed to exert their activity through the upregulation of neurotrophic factor synthesis. The upregulation of neurotrophic factor synthesis can involve one or more of these neurotrophic factors: NGF, NT-3, BDNF, and NT-4/5.

Exemplary studies and treatments were performed as discussed below using
 30 various dosages and routes of administration of selected exemplary compounds representative of compositions that are effective with the methods of the present invention. Of course, those skilled in the art will recognize that the present invention is

not specifically limited to the particular compositions, dosages or routes of administration detailed below.

Depending upon the particular needs of the individual subject involved, the compositions used in the present invention may be administered in various doses to provide effective treatment concentrations based upon the teachings of the present invention. What constitutes an effective amount of the selected composition will vary based upon such factors including the activity of the selected compound, the physiological characteristics of the subject, the extent and nature of the subject's disease or condition and the method of administration. Exemplary treatment concentrations which have proven effective in modifying neural activity range from less than 1 μM to concentrations of 500 mM or more. Generally, initial doses will be modified to determine the optimum dosage for treatment of the particular mammalian subject. The compositions may be administered using a number of different routes including orally, topically, transdermally, intraperitoneal injection or intravenous injection directly into the bloodstream. Of course, effective amounts of the compounds may also be administered through injection into the cerebrospinal fluid or infusion directly into the brain, if desired.

The methods of the present invention may be effected using compounds administered to a mammalian subject either alone or in combination as a pharmaceutical formulation. Further, the compounds may be combined with pharmaceutically acceptable excipients and carrier materials such as inert solid diluents, aqueous solutions or non-toxic organic solvents. If desired, these pharmaceutical formulations may also contain preservatives and stabilizing agents and the like, as well as minor amounts of auxiliary substances such as wetting or emulsifying agents, as well as pH buffering agents and the like which enhance the effectiveness of the active ingredient. The pharmaceutically acceptable carrier can be chosen from those generally known in the art, including, but not limited to, human serum albumin, ion exchangers, dextrose, alumina, lecithin, buffer substances such as phosphate, glycine, sorbic acid, potassium sorbate, propylene glycol, polyethylene glycol, and salts or electrolytes such as protamine sulfate, sodium chloride, or potassium chloride. Other carriers can be used.

Liquid compositions can also contain liquid phases either in addition to or to the exclusion of water. Examples of such additional liquid phases are glycerin, vegetable

oils such as cottonseed oil, organic esters such as ethyl oleate, and water-oil emulsions.

The compositions can be made into aerosol formations (i.e., they can be "nebulized") to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichloromethane, propane, or nitrogen. Other suitable propellants are known in the art.

Formulations suitable for parenteral administration, such as, for example, by intravenous, intramuscular, intradermal, and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions. These can contain antioxidants, buffers, preservatives, bacteriostatic agents, and solutes that render the formulation isotonic with the blood of the particular recipient. Alternatively, these formulations can be aqueous or non-aqueous sterile suspensions that can include suspending agents, thickening agents, solubilizers, stabilizers, and preservatives. Compositions suitable for use in methods according to the present invention can be administered, for example, by intravenous infusion, orally, topically, intraperitoneally, intravesically, or intrathecally. Formulations of compounds suitable for use in methods according to the present invention can be presented in unit-dose or multi-dose sealed containers, in physical forms such as ampules or vials.

The drug-induced peripheral neuropathy to be treated can be drug-induced peripheral neuropathy induced by the administration of a vinca alkaloid, cisplatin, paclitaxel, suramin, altretamine, carboplatin, chlorambucil, cytarabine, dacarbazine, docetaxel, etoposide, fludarabine, ifosfamide with mesna, tamoxifen, teniposide, and thioguanine. Vinca alkaloids include vincristine and vinblastine. Peripheral neuropathy associated with the administration of oncolytic drugs is described in C.M. Haskell, "Cancer Treatment" (5th Ed., W.B. Saunders, Philadelphia, 2001), ch. 10, pp. 104-214, incorporated herein by this reference. In particular, methods according to the present invention are particularly useful in treating drug-induced peripheral neuropathy associated with the administration of vincristine, paclitaxel, or cisplatin.

Although Applicants do not intend to be bound by this theory, the beneficial effects of bifunctional purine derivatives such as AIT-082 may depend on generalized trophic and NGF-sensitive mechanisms and not merely on the induction of sprouting. These trophic effects may boost the capacity of NGF-sensitive neurons to respond to still unknown regeneration factors other than NGF itself. For example, it is likely that

CGRP and Substance P expression increases with treatment with AIT-082 or other bifunctional purine derivatives.

The invention is illustrated by the following Examples. These Examples are presented for illustration only and are not intended to limit the invention.

5

EXAMPLE 1

EFFECT OF AIT-082 ON NEUROTOXICITY INDUCED BY CISPLATIN AND

VINCRIStINE

METHODS AND MATERIALS

In Vitro Procedures

10

Culture of dorsal root ganglion neurons.

The dorsal root ganglion neurons were obtained from rat fetuses (Wistar rats; E15 ; Elevage Janvier, Le Genest St Isle). They were dissected and incubated in trypsin 1X (Gibco; Life Technologies, Cergy-Pontoise, France) for 30 min at 37°C. The reaction was stopped by addition of free Hanks balanced salt solution containing calcium and magnesium ; (HBSS⁺ ; Gibco) added with 10 % of fetal bovine serum (FBS ; Gibco) and 0.5 mg/ml DNase I (Boehringer Mannheim, Meylan, France). The suspension was triturated with a 10-ml pipette and using a needle syringe. After centrifugation, the dissociated cells were resuspended in DMEM containing 10 % FBS, 2% of chick embryos extract (Gibco), 1 % antibiotics (Gibco) and NGF (Tébu, Le Perray en Yvelines, France) 3 ng/ml final.

20

Viable cells were counted and seeded at $\pm 50,000$ cells /well in 96 well-plates (Nunc, Life Technologies) coated with poly-L-lysine (0.01 mg/ml, Sigma). Cells were maintained in a humidified incubator at 37°C in 5 % CO₂-95 % air atmosphere.

After 24h, the culture medium was completed with anti-mitotic compounds (5-fluoro-5'-deoxyuridine and cytosine β -D-arabinofuranoside) and uridine all 10⁻⁵M final. This treatment is aimed at eliminating contaminating fibroblasts and inhibiting the proliferation of Schwann cells.

25

Cultures were rinsed 48 hrs later (medium without any growth factors) and medium was replaced with fresh medium (200 μ l /well) containing cisplatin (3.5 ng/ml ; Sigma) added or not with the reference or test compounds.

30

After 48 h of intoxication, survival rates were measured using an acid phosphatase enzymatic activity assay.

Survival rate.

Acid phosphatase activity was measured according to a previously described method (Ueda et al.; 1994). After removal of the culture medium, wells were rinsed twice with PBS (Life Technologies) and incubated with 100 µl of buffer containing 0.1 M sodium acetate (pH5.5), 0.1 % Triton X100 (Sigma) and 10 mM p-nitrophenyl phosphate (Sigma) 1h at 37°C/5% CO₂. Reaction was stopped by addition of 10 µl of 1N NaOH (LPCR, Strasbourg, France). Enzyme activity was measured at 405 nm in a microplate reader (LabSystem, France).

Acid phosphatase activity is proportional to the amount of living cells. The results were expressed as percentage O.D. as compared to control conditions (same rinsing and medium replacements, but no intoxication and no treatment with either NGF, riluzol or the test compounds).

In Vivo Procedures

Animals received vincristine 0.15 mg/ml (IP) per day with the following timing: 5 days of intoxication followed with 2 days without vincristine injection and a new period of 5 days of intox followed with 2 days no intoxication and 2 other days of intoxication with vincristine followed by a 3 week recovery period. The test compound was administered daily from the first day of vincristine injection to the end of the experiments

Subjects used were 65 female Dark Agouti rats (200-250 g) (n = 13 per group). Five groups were used: (1) a vehicle control group; (2) a vincristine-vehicle group; (3) a vincristine + AIT-082 3 mg/kg IP group; (4) a vincristine + AIT-082 10 mg/kg IP group; and (5) a vincristine + AIT-082 30 mg/kg IP group.

Behavioral measurements. The general health and weight of animals were checked every day. Sensorimotor tests were performed twice a week during 6 week study, starting one week before beginning of intoxication (baseline).

Electrophysiological measurements. Electrophysiological recordings were performed once a week using a Neuromatic 2000M electromyographical apparatus (Dantec, France). EMG were performed under Ketamine anesthesia (Imalgène 500®, Merieux, France).

The measured parameters were: the sensitive nerve conduction velocity (SNCV), the amplitude and the latency compound muscle action potential (CMAP).

Sensitive nerve conduction velocity. Skin electrodes were used. Caudal nerve was stimulated with a series of 20 pulses during 0.2 ms at a supramaximal intensity. The mean of 20 stimulations is included for statistical analysis.

Compound muscle action potential and distal latency. CPAM was measured in gastrocnemius muscle after stimulation of the sciatic nerve. Sciatic nerve was stimulated with a single 0.2 ms pulse at a supramaximal intensity (12.8 mA). The amplitude (mV) and the latency of the response (ms) are measured. The amplitude is indicative of the number of active motor units, while distal latency reflects motor nerve conduction velocity.

Drug administration. The test compound was administered daily by the i.p. route from the first day of intoxication with the vincristine to the end of the experiments.

Morphometric analysis. The morphometric analysis was performed on 3 animals per group on sciatic nerve. It includes the measurements of various parameters according to the following procedure:

(1) Dissection of a segment of sciatic nerve (2 cm from the spinal cord, L5-L6, to the trifurcation of the sciatic).

The tissue was fixed overnight with glutaraldehyde 4% in phosphate buffer (PH 7.4) and then maintained in 30% sucrose in phosphate buffer at + 4°C until use. The sciatic nerve was fixed in 2% osmium tetroxide in phosphate buffer for 2 hours, dehydrated in serial alcohol solutions and embedded in Eppon. Embedded tissues were then placed at +70°C for 3 days.

(2) Microtome sections : Transverse sections of 1.5 µm were made with a microtome and stained with toluidine blue, then dehydrated and mounted in Eurokitt.

(3) Morphometric analysis : Sections were observed using an optical microscope (Nikon) and morphometric analysis were performed with a semi-automated digital image analysis software (Morphonorf, Alcatel). Per slice, 4 fields were analyzed, the number of myelinated fibers, density of fibers, diameter of fibers, thickness of myelin sheath, surface of myelin sheath and G factor.

RESULTS

In Vitro Results. AIT-082 at a concentration of 1 µM significantly increased survival compared to cisplatin alone (Figure 1). These effects diminished progressively with higher doses. Neotrofin was 10 times more potent than riluzole and had a greater effect than 5 ng/ml of NGF. AIT-034 was ineffective at all concentrations (0.01-100 µM).

In Vivo Results. Vincristine treatments (alone and with AIT-082) produced significant reductions in body weight with repetitive dosing compared to day matched

controls with vehicle treatment (Figure 2) at days 11-19 and 22-31. These effects all these treatments were reversible by the termination of treatments. This suggests that the dosage of vincristine and the route of administration did not induce permanent deterioration of health. AIT-082 in combination with vincristine produced a pattern of dose-related reductions in body weight below that observed with control or vincristine alone. The body weights of the animals treated with AIT-082 at 30 mg/kg were significantly lower than treatment with vincristine alone at days 11, 14, 16, 18, 24 and 25. These effects were also reversible and did not appear to effect the improvements in sensory nerve conduction or EMG amplitude described below.

AIT-082 produced significant and dose-related changes in sensory nerve conduction velocity at day 17 during the initial phase of intoxication to vincristine and day 31 during the recovery phase of the study. These finding are suggestive that AIT-082 can lessen the severity of the neuropathy and increase the recovery to normal condition.

AIT-082 produced significant and dose-related changes in EMG amplitude at days 17 and 23 during the initial phase of intoxication to vincristine and days 31 and 38 during the recovery phase of the study. These finding are suggestive that AIT-082 can lessen the severity of the neuropathy and improve recovery. Although these data can not distinguish between effects on muscle responses and effects on nerve conduction, they are supportive of the findings on sensory nerve conduction velocity.

At the end of the treatment periods, AIT-082 treatment was found to have produced greater numbers of myelinated nerve fibers in sciatic nerve biopsies than in animals treated with vincristine alone. These findings are also supportive of either a preventative or regenerative effect of AIT-082 on neurotoxicity induced by vincristine.

DISCUSSION

In an in vitro model for neurotoxicity induced by oncolytic drugs, AIT-082 produced statistically significant protection against cisplatin induced neurotoxicity. These findings indicate that AIT-082 given simultaneously to with an oncolytic drug in patients undergoing cancer therapy could prevent the neuropathic side-effects. Further confirmation of this was obtained from an in vivo model for neurotoxicity induced by oncolytic drugs. In this model, AIT-082 treatment reduced the severity the vincristine induced changes in sensory nerve conduction velocity and electromyographic responses and improved the rate of recovery after the termination of vincristine therapy. The in vivo studies are supportive of a potential use of AIT-082

and related compounds for the prevention of drug-induced neuropathies and also suggest that these compounds could be used to improve recovery from these neuropathies.

Clinical implications of the present findings.

5 There is increasing evidence that deficient neurotrophic support, including that provided to cutaneous nerves by NGF, contributes to the pathogenesis of drug-induced neuropathies, such as oncolytic therapies. In animal studies, Hayakawa et al. 1998 demonstrated that NGF could prevent neurotoxicity induced by vincristine, taxol and cisplatin. In clinical studies, patients suffering from neurological deficits related to
10 treatments with antitumor agents have low circulating levels of NGF (De Santis et al., 2000).

REFERENCES

K. Hayakawa, T. Itoh, H. Niwa, T. Mutoh and G. Sobue, "NGF prevention of neurotoxicity induced by cisplatin, vincristine and taxol depends on toxicity of each
15 drug and NGF treatment schedule: In vitro study of adult rat sympathetic ganglion explants. Brain Research 794:313-319 (1998)

S. De Santis, A. Pace, L. Bove, F. Cognetti, F. Properzi, M. Fiore, V. Triaca, A. Savarese, M. D. Simone, B. Jandolo, L. Manzione and I Aloe. "Patients treated with antitumor drugs displaying neurological deficits are characterized by a low circulating
20 level of Nerve Growth Factor." Clinical Cancer Research 6: 90-95, 2000.

Further, indirect support for this therapeutic approach comes from our earlier findings (Diamond et al., 1988; 1992a) that chronic NGF-deprivation causes a shrinkage of nociceptive fields in the skin consistent with a "dying-back" neuropathy. AIT-082 administration, were it to induce endogenous NGF increases in the skin of
25 patients taking oncolytic drugs, could help protect NGF-sensitive neurons from the threat of neuropathy, without the hazard of hyperalgesia, as explained above.

REFERENCES

The following references are cited in Example 1:

Albers KM, Wright DE, Davis BM (1994) Overexpression of nerve growth factor
30 in epidermis of transgenic mice causes hypertrophy of the peripheral nervous system. J Neurosci 14:1422-1432.

Campenot RB (1994) NGF and the local control of nerve terminal growth. J Neurobiol 25:599-611

Dantes M and McComas A (1981) The extent and time course of motoneuron involvement in amyotrophic lateral sclerosis. *Muscle and Nerve* 14:416-421.

Davis BM, Fundin BT, Albers KM, Goodness TP, Cronk KM, Rice FL (1997) Overexpression of nerve growth factor in skin causes preferential increases among
5 innervation to specific sensory targets. *J Comp Neurol* 387:489-506

Diamond J, Cooper G and Turner C (1976) Trophic regulation of nerve sprouting. *Science* 193:371-377.

Diamond J, Holmes M and Visheau B (1988) NGF-regulated plasticity in the adult nervous system. *Soc Neurosci Abstr* 14:245.6

10 Diamond J, Coughlin M, MacIntyre L, Holmes M and Visheau B (1987) Evidence that endogenous nerve growth factor is responsible for the collateral sprouting, but not regeneration, of nociceptive axons in adult rats. *Proc Natl Acad Sci USA* 84:6596-6600.

Diamond J, Coughlin M and Holmes M (1992a) Endogenous NGF and impulses
15 regulate the collateral sprouting of sensory nerves in the skin of the adult rat. *J. Neurosci* 12:1454-1466.

Diamond J, Holmes M, Foerster A and Coughlin M (1992b) Sensory nerves in adult rats regenerate and restore sensory function to the skin independently of endogenous NGF. *J. Neurosci* 12:1467-1476.

20 Doucette R, Diamond J (1987) The normal and precocious sprouting of heat nociceptors in the skin of adult rats. *J. Comp. Neurol* 261:592-603.

English KB, Harper S, Stayner N, Wang ZM, Davies AM (1994) Localization of nerve growth factor (NGF) and low-affinity NGF receptors in touch domes and quantification of NGF mRNA in keratinocytes of adult rats. *J. Comp. Neurol* 344:470-
25 480.

Gloster A, Diamond J (1992) Sympathetic nerves in adult rats regenerate normally and restore pilomotor function during an anti-NGF treatment that prevents their collateral sprouting. *J. Comp. Neurol* 326:363-374.

Imayama S (1981) Scanning and transmission electron microscope study on
30 the terminal blood vessels of the rat skin. *J. Invest Dermatol* 76:151-157.

Jackson PC, Diamond J (1984) Temporal and spatial constraints on the collateral sprouting of low-threshold mechanosensory nerves in the skin of rats. *J. Comp. Neurol* 226:336-345.

- Karchewski LA, Kim FA, Johnston J, McKnight RM, Verge VM (1999) Anatomical evidence supporting the potential for modulation by multiple neurotrophins in the majority of adult lumbar sensory neurons. *J Comp Neurol* 413(2):327-41.
- 5 Korsching S, Theonen H (1985) Nerve growth factor supply for sensory neurons: site of origin and competition with the sympathetic nervous system. *Neurosci Lett* 54:201-205.
- Levi-Montalcini R, Skaper SD, Dal Toso R, Petrelli L, Leon A. 1996 Nerve growth factor: from neurotrophin to neurokine. *TINS* 19:514-520.
- 10 Lewin GR, Ritter AM, Mendell LM (1993) Nerve growth factor-induced hyperalgesia in the neonatal and adult rat. *J. Neurosci* 13:2136-2148.
- Lewin GR, Rueff A, Mendell LM (1994) Peripheral and central mechanisms of NGF-induced hyperalgesia. *Eur J Neurosci* 6:1903-1912.
- Mearow KM, Kril Y, and Diamond J (1993) Increased NGF mRNA expression in denervated rat skin. *Neuroreport* 4:351-354.
- 15 Murphy RA, Acheson A, Hodges R, Haskins J, Richards C, Reklow E V, Chlumecky V, Barker PA, Alderson RF, Lindsay RM (1993) Immunological relationships of NGF, BDNF and NT3; recognition and functional inhibition by antibodies to NGF. *J. Neurosci* 13:2853-2862.
- 20 Neumann S, Doubell TP, Leslie T, Woolf CJ. 1996 Inflammatory pain hypersensitivity mediated by phenotypic switch in myelinated primary sensory neurons. *Nature* 384:360-364.
- Nixon BJ, Doucette R, Jackson P and Diamond J (1984) Impulse activity evokes precocious sprouting of nociceptive nerves into denervated skin. *Somatosensory Res* 2:97-126.
- 25 Pertens E, Urschel-Gybers BA, Holmes M, Pal R, Foerster A, Kril Y and Diamond J (1999) Intraspinal and behavioural consequences of NGF-induced nociceptive sprouting and NGF induced hyperalgesia compared in adult rats. *J Comp Neurol* 410:73-89.
- 30 Ramer MS, Kawaja MD, Henderson JT, Roder JC, Bisby MA (1998) Glial overexpression of NGF enhances neuropathic pain and adrenergic sprouting into DRG following sciatic nerve constriction in mice. *Neurosci Lett* 251 53-56
- Romero MI, Rangappa N, Li L, Lightfoot E, Garry MG, Smith GM (2000) Extensive sprouting of sensory afferents and hyperalgesia induced by conditional expression of nerve growth factor in the adult spinal cord. *J Neurosci* 20:4435-4445

Stucky CL, Koltzenburg M, Schneider M, Engle MG, Albers KM, Davis BM (1999) Overexpression of nerve growth factor in skin selectively affects the survival and functional properties of nociceptors. *J Neurosci* 19:8509-8516

Theriault E, Diamond J (1988) Nociceptive cutaneous stimuli evoke localized
5 contractions in a skeletal muscle. *J Neurophysiol* 60:446-462.

Toma JG, Rogers D, Senger DI, Campenot RB, Miller FD (1997) Spatial regulation of neuronal gene expression in response to nerve growth factor. *Dev Biol* 184: 1-9

Yasargil GM, Macintyre L, Doucette R, Visheau B, Holmes M and Diamond J
10 (1988) Axonal domains within shared touch domes in the rat: a comparison of their fate during conditions favoring collateral sprouting and following axonal regeneration. *J Comp Neurol* 270:301-312.

EXAMPLE 2

FURTHER NEUROPROTECTIVE EFFECTS OF AIT-082

15 The objectives of the studies described in this application are to evaluate AIT-082 and related analogues for their potential utility in the treatment of neuropathies associated with oncolytic therapies. The prototypic compound, AIT-082, is currently in Phase 2 clinical trials for Alzheimer's disease (AD) and has been proven safe in over 1400 patients. Its preclinical pharmacological effects include neuroprotection against
20 excitotoxicity, induction of the synthesis and secretion of neurotrophic and pleiotrophic factors, neuroregenerative effects and proliferative effects on neural progenitor cells. The effects of AIT-082 are unlike any product that has been approved or is currently under evaluation for neuropathies associated with oncolytic agents. AIT-082 and its second generation analogues may arrest or retard neurodegenerative processes,
25 promote neuroregeneration and provide symptomatic relief for cancer patients with neuropathies or at high risk of developing neuropathies.

Background: AIT-082 (Neotrofin™, leteprinim potassium, 4[[3(1,6-dihydro-6-oxo-9-purin-9-yl)-1-oxopropyl] amino] benzoic acid, potassium salt) is a novel
30 therapeutic agent currently in clinical testing for the treatment of AD. The proposed preclinical studies and clinical trials for AIT-082 in oncolytic therapy-induced neuropathies are based upon its neuroprotective, neurotrophic and neuroregenerative effects in preclinical research. These effects may prevent or reverse the progressive degenerative changes associated with oncolytic therapies.

In vitro effects on expression and secretion of nerve growth factors:

Modulation of the secretion or efficacy of neurotrophic factors may have a major role in preventing or reversing neuropathies due to oncolytic therapies^{3,4}. Findings from studies with rat astrocyte cultures provided evidence that AIT-082 increases the production of mRNAs for several neurotrophic factors, including NGF, basic fibroblast growth factor (bFGF), and neurotrophin 3 (NT-3)^{1,2}. In astrocytes, AIT-082 enhanced the secretion into the culture medium of the neurotrophic proteins NGF, TGF and S 100⁵. The secretion of NGF and TGF were dependent upon *de novo* protein synthesis induced by AIT-082 as indicated by the suppression of these effects by cycloheximide pre-treatment. The secretion of S 100 was independent of protein synthesis and from existing cellular stores after AIT-082 treatment.

In vivo effects on expression and secretion of nerve growth factors: In

attempt to further explore the effects of AIT-082 that could be beneficial for chemotherapy-induced neuropathy, additional studies were conducted on dose-related effects on BDNF levels in the spinal cord. In these studies, AIT-082 was administered in drinking water for 7 days and BDNF content was measured by ELISA assay. AIT-082 produced significant dose-related increases in BDNF levels in the rat spinal cord.

The following are the references for Example 2:

1. Glasky AJ, Glasky MS, Ritzmann RF and Rathbone MP. AIT-082, a novel purine derivative with neuroregenerative properties. *Exp. Opin. Invest. Drugs.* 6:1413-1417, 1997.
2. Rathbone MP, Middlemiss PJ, Gysbers J, Diamond J, Holmes M, Pertens E, Juurlink BH, Glasky A, Ritzmann R, Glasky M, Crocker CE, Ramirez JJ, Lorenzen A, Fein T, Schultze E, Schwabe U, Ciccarelli R, Di Iorio P, and Caciagli F. Physiology and pharmacology of natural and synthetic nonadenine-based purines in the nervous system. *Drug. Devel. Res.* 45:356-372, 1998.
3. Hayakawa K, Sobue G, Itoh T. Nerve growth factor prevents neurotoxic effects of cisplatin, vincristine and taxol, on adult rat sympathetic ganglion explants in vitro. *Life Sci* 55:591-525, 1994.
4. Hayakawa K, Itoh T, Niwa H, Mutoh T, Sobue G, NGF prevention of neurotoxicity induced by cisplatin, vincristine and taxol depends on toxicity of each drug and NGF treatment schedule: in vitro study of adult rat sympathetic ganglion explants. *Brain Res* 794:313-319, 1998.

5. Middlemiss P, Glasky AJ, Rathbone MP, Werstuik E, Hindley S and Gysbers J. AIT-082, a unique purine derivative, enhances nerve growth factor mediated outgrowth from PC12 cells. *Neurosci. Letters* 199:1-4, 1995.

ADVANTAGES OF THE INVENTION


5 The present invention provides new methods for treating patients with drug-induced peripheral neuropathy, including peripheral neuropathy associated with the administration of oncolytic drugs, to induce peripheral nerve sprouting, which can include nociceptive nerve sprouting. These methods provide for nerve regeneration. These methods can be performed, at least in some alternatives, without inducing
10 hyperalgesia. These methods can be combined with other treatments for drug-induced peripheral neuropathy, such as palliative measures for the relief of pain. They do not depend on specific interactions between the compounds administered and the

drugs causing peripheral neuropathy.


Although the present invention has been described in considerable detail, with
15 reference to certain preferred versions thereof, other versions and embodiments are possible. Therefore, the scope of the invention is determined by the following claims.

WE CLAIM:

1. A method of treating drug-induced neuropathy comprising administering to a patient with drug-induced neuropathy an effective amount of a compound having activity against drug-induced neuropathy, the compound comprising: (1) a moiety A selected from the group consisting of a purine moiety, a purine analogue, a tetrahydroindolone moiety, a tetrahydroindolone analogue, a pyrimidine moiety, and a pyrimidine analogue; (2) a hydrocarbyl moiety L of 1 to 6 carbon atoms that is linked to the moiety A and that can be cyclic, with the hydrocarbyl moiety being optionally substituted with one or more substituents selected from the group consisting of lower alkyl, amino, hydroxy, lower alkoxy, lower alkylamino, lower alkylthio, and oxo; and (3) a moiety B that is linked to the moiety L wherein B is $-OZ$ or $N(Y_1)-D$, where Z is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, aralkyl, or heteroaralkyl; D is a moiety that promotes absorption of the compound having activity against drug-induced neuropathy; and Y_1 is hydrogen, alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms, which can be N, O, or S.
2. The method of claim 1 wherein the compound having activity against drug-induced neuropathy passes through the blood-brain barrier.
3. The method of claim 1 wherein A is a purine moiety.
4. The method of claim 3 wherein A is a substituted or unsubstituted hypoxanthine moiety.
5. The method of claim 4 wherein L has the structure $-(CH_2)_n-CONH-$ where n is an integer from 1 to 6.
6. The method of claim 5 wherein the compound having activity against drug-induced neuropathy is a compound of formula (I)



where n is an integer from 1 to 6 and R is hydrogen or lower alkyl or is a salt or prodrug ester of a compound of formula (I)



wherein n is an integer from 1 to 6 and R is hydrogen or lower alkyl.

7. The method of claim 6 wherein the compound having activity against drug-induced neuropathy is a compound of formula (I) wherein n is an integer from 1 to 6 and R is hydrogen or lower alkyl.


8. The method of claim 7 wherein R is hydrogen.

9. The method of claim 8 wherein n is 2 and the compound is N-4-[[3-(1,6-dihydro-6-oxo-purin-9-yl)-1-oxopropyl] amino] benzoic acid.

10. The method of claim 7 wherein R is ethyl.

11. The method of claim 10 wherein n is 2 and the compound is N-4-[[3-(1,6-dihydro-6-oxo-purin-9-yl)-1-oxopropyl] amino] benzoic acid ethyl ester.


12. The method of claim 5 wherein the compound having activity against drug-induced neuropathy is a compound of formula (II)



wherein n is an integer from 1 to 6, R is selected from the group consisting of H, $COOH$, and $COOW_1$, wherein W_1 is selected from the group consisting of lower alkyl, amino, and lower alkylamino, and R_2 is selected from the group consisting of H and OH.

13. The method of claim 12 wherein n is 2.

14. The method of claim 5 wherein the compound having activity against drug-induced neuropathy is a compound of formula (III)




wherein n is an integer from 1 to 6, R_1 is selected from the group consisting of H, $COOH$, and $COOW_1$, wherein W_1 is selected from the group consisting of lower alkyl, amino, and lower alkylamino, R_2 is selected from the group consisting of H and OH, and R_3 is selected from the group consisting from the group consisting of H and OH.

15. The method of claim 14 wherein n is 2.

16. The method of claim 3 wherein A is a substituted or unsubstituted guanine moiety.

17. The method of claim 16 wherein L has the structure $-(CH_2)_n-CONH-$ wherein n is an integer from 1 to 6.

18. The method of claim 17 wherein the compound having activity against drug-induced neuropathy is a compound of formula (IV)




wherein n is an integer from 1 to 6, R_1 is selected from the group consisting of H, COOH, and COOW₁, wherein W₁ is selected from the group consisting of lower alkyl, amino, and lower alkylamino and R_2 is selected from the group consisting of H and OH.

19. The method of claim 18 wherein n is 2, R_1 is H, and R_2 is OH, and the compound is N-(2-(5-hydroxyindol-3-yl)) ethyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

20. The method of claim 18 wherein n is 2, R_1 is H, and R_2 is H, and the compound is N-(2-(2-indol-3-yl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

21. The method of claim 18 wherein n is 2, R_1 is COOH, and R_2 is OH, and the compound is N-(1-carboxyl-(2-(5-hydroxyindol-3-yl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

22. The method of claim 17 wherein the compound having activity against drug-induced neuropathy is a compound of formula (V)




wherein n is an integer from 1 to 6 and R is selected from the group consisting of hydrogen and lower alkyl.

23. The method of claim 22 wherein n is 2, R is hydrogen, and the compound is N-4-carboxyphenyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

24. The method of claim 22 wherein n is 2, R is ethyl, and the compound is N-4-carboxyphenyl-3-(2-amino-6-oxohydropurin-9-yl) propanamide ethyl ester.

25. The method of claim 17 wherein the compound having activity against drug-induced neuropathy is a compound of formula (VI)




wherein n is an integer from 1 to 6 and R is selected from the group consisting of hydrogen and lower alkyl.

26. The method of claim 25 wherein n is 2, R is hydrogen, and the compound is 3-(2-amino-6-oxohydropurin-9-yl) propanoic acid.

27. The method of claim 25 wherein n is 2, R is ethyl, and the compound is 3-(2-amino-6-oxohydropurin-9-yl) propanoic acid ethyl ester.


28. The method of claim 17 wherein the compound having activity against drug-induced neuropathy is a compound of formula (VII)



wherein n is an integer from 1 to 6, p is an integer from 1 to 6, and q is an integer from 1 to 3.

29. The method of claim 28 wherein n is 2, p is 2, and q is 1, and the compound is N-[2-[[2-(2-oxopyrrolidin-1-yl)-1-oxoethyl] amino] ethyl] propanamide.

30. The method of claim 17 wherein the compound having activity against drug-induced neuropathy is a compound of formula (VIII)




wherein n is an integer from 1 to 6, R_1 is selected from the group consisting of H, COOH, and COOW₁, wherein W₁ is selected from the group consisting of lower alkyl, amino, and lower alkylamino, R_2 is selected from the group consisting of H and OH, and R_3 is selected from the group consisting of H and OH.

31. The method of claim 30 wherein n is 2, R_1 is H, R_2 is H, and R_3 is OH, and the compound is N-(2-(3,4-dihydroxyphenyl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

32. The method of claim 30 wherein n is 2, R_1 is H, R_2 is OH, and R_3 is OH, and the compound is N-(2-hydroxy-2-(3,4-dihydroxyphenyl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

33. The method of claim 30 wherein n is 2, R_1 is COOH, R_2 is H, and R_3 is H, and the compound is N-(1-carboxyl-2-(3,4-dihydroxyphenyl)ethyl)-3-(2-amino-6-oxohydropurin-9-yl) propanamide.

34. The method of claim 16 wherein the compound having activity against drug-induced neuropathy is a compound of formula (IX)




wherein n is an integer from 1 to 6 and p is an integer from 1 to 3.

35. The method of claim 34 wherein n is 2, p is 1, and the compound is N-4-[[3-(2-amino-6-oxohydropurin-9-yl) 1-oxopropyl] amino] benzoic acid 1-(dimethylamino)-2-propyl ester.

36. The method of claim 1 wherein A is a substituted or unsubstituted 9-atom bicyclic moiety in which the 5-membered ring has 1 to 3 nitrogen atoms, the

bicyclic moiety having the structure of formula (X)



where:

(a) if the bond between N₁ and the bond between C₅ is a single bond, then the bond between C₆ and R₆ is a double bond, R₆ is O or S, and R₁ is hydrogen, alkyl, aralkyl, cycloalkyl, or heteroaralkyl;

(b) if the bond between N₁ and C₆ is a double bond, then the bond between C₆ and R₆ is a single bond, R₁ is not present, and R₆ is hydrogen, halo, amino, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁, where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(c) if the bond between C₂ and N₃ is a single bond, then the bond between C₂ and R₂ is a double bond, R₂ is O or S, and R₃ is hydrogen or alkyl;

(d) if the bond between C₂ and N₃ is a double bond, then the bond between C₂ is a single bond, R₃ is not present, and R₂ is hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, amino, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁,

where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, aralkoxy carbonyl, heteroaralkoxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(e) A₇ and A₈ are C or N;

(i) if A₇ and A₈ are both C and the bond between A₇ and A₈ is a single bond, then the bond between A₈ and R₈ is two single bonds to two hydrogen atoms or is a double bond in which R₈ is O or S and R₇ is two hydrogen atoms;

(ii) if A₇ and A₈ are both C and the bond between A₇ and A₈ is a double bond, then R₇ is hydrogen, the bond between A₈ and R₈ is a single bond and R₈ is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

(iii) if A₇ and A₈ are both N, then the bond between A₇ and A₈ is a double bond, and R₇ and R₈ are not present;


(iv) if A₇ is C and A₈ is N, then the bond between A₇ and A₈ is a double bond, R₇ is hydrogen, and R₈ is not present;

(v) if A₇ is N, A₈ is C, and the bond between A₇ and A₈ is a double bond, then R₇ is not present, the bond between A₈ is a single bond, and R₈ is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

(vi) if A₇ is N, A₈ is C, and the bond between A₇ and A₈ is a single bond, then R₇ is hydrogen, alkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl, the bond between A₈ and R₈ is a double bond, and R₈ is O or S; and

(f) N_9 is bonded to L ; with the proviso that A does not have the structure of an unsubstituted guanine or hypoxanthine.

37. The method of claim 3 wherein the purine moiety is a purine moiety of formula (XI)



in which:

(a) R_1 is selected from the group consisting of hydrogen, alkyl, aralkyl, cycloalkyl, and heteroaralkyl; and R_2 is selected from the group consisting of hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, OQ_1 , SQ_1 , $NHNH_2$, $NHOQ_1$, NQ_1Q_2 , or NHQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_1 and Q_2 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, arylkoxycarbonyl, heteroarylokoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroarylalkylaminocarbonyl in which the alkyl portions could be cyclic and can contain from one to three heteroatoms which could be N, O, or S, with the proviso that both R_1 and R_2 are not hydrogen and that R_1 is not hydrogen when R_2 is amino.

38. The method of claim 37 wherein R_1 is butyl and R_2 is hydrogen.


39. The method of claim 37 wherein R_1 is benzyl and R_2 is hydrogen.

40. The method of claim 37 wherein R_1 is dimethylaminoethyl and R_2 is hydrogen.

41. The method of claim 37 wherein R_1 is cyclopentyl and R_2 is hydrogen.

42. The method of claim 37 wherein R₁ is cyclohexylmethyl and R₂ is hydrogen.
43. The method of claim 37 wherein R₁ is cyclopropylmethyl and R₂ is hydrogen.
44. The method of claim 37 wherein R₁ is hydrogen and R₂ is phenyl.
45. The method of claim 37 wherein R₁ is hydrogen and R₂ is butyl.
46. The method of claim 37 wherein R₁ is butyl and R₂ is butyl.
47. The method of claim 37 wherein R₁ is hydrogen and R₂ is methyl.
48. The method of claim 37 wherein R₁ is hydrogen and R₂ is phenylamino.
49. The method of claim 3 wherein the purine moiety is a purine moiety of

Formula (XII)



in which:

(a) R₂ is selected from the group consisting of hydrogen, halo, amino, OQ₃, SQ₃, NHNH₂, NHOQ₃, NQ₃Q₄, or NHQ₃, where Q₃ and Q₄ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, and heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₃ and Q₄ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₃ where Y₃ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaryloxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and

(b) R_6 is selected from the group consisting of hydrogen, halo, amino, OQ_5 , SQ_5 , $NHNH_2$, $NHOQ_5$, NQ_5Q_6 , or NHQ_6 , where Q_5 and Q_6 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, and heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_5 and Q_6 are present together and are alkyl, they can be taken together to form a 5- or 6- membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, arylkoxycarbonyl, heteroarylkoxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S.

50. The method of claim 49 wherein R_2 is hydrogen and R_6 is amino.


51. The method of claim 49 wherein R_6 is chloro.

52. The method of claim 49 wherein R_6 is phenylamino.

53. The method of claim 49 wherein R_2 is amino and R_6 is chloro.

54. The method of claim 3 wherein the purine moiety is the purine moiety of

Formula (XIII)



in which:

(a) R_1 is hydrogen, alkyl, aralkyl, cycloalkyl, or heteroaralkyl; and

(b) R_2 is O or S.

55. The method of claim 54 wherein R_1 is hydrogen.


56. The method of claim 54 wherein R_2 is O.

57. The method of claim 54 wherein R_2 is S.
58. The method of claim 3 wherein the compound is 4-[3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester.
59. The method of claim 3 wherein the compound is 4-[3-(1-butyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester.
60. The method of claim 3 wherein the compound is 4-[3-(1-methyl-6-oxo-1,6-dihydropurin-9-yl)propionylamino] benzoic acid ethyl ester.
61. The method of claim 3 wherein the compound is 4-[3-(1,2-dimethylaminoethyl)-6-oxo-1,6-dihydropurin-9-yl) propionylamino] benzoic acid ethyl ester.
62. The method of claim 3 wherein the compound is 4-[3-(2,6-dioxo-1,2,3,6-tetrahydropurin-9-yl) propionylamino] benzoic acid ethyl ester.
-
63. The method of claim 3 wherein the compound is 4-[3-(6-methoxypurin-9-yl) propionylamino] benzoic acid ethyl ester.
64. The method of claim 3 wherein the compound is 4-[3-(6-dimethylaminopurin-9-yl) propionylamino] benzoic acid ethyl ester.
65. The method of claim 3 wherein the compound is 4-[3-(2-amino-6-chloropurin-9-yl) propionylamino] benzoic acid ethyl ester.
66. The method of claim 3 wherein the compound is 4-[2-(6-oxo-2-thioxo-1,2,3,6-tetrahydropurin-9-yl) propionylamino] benzoic acid ethyl ester.
67. The method of claim 3 wherein the compound is 4-[2-(2-butyl-6-oxo-1,6-dihydropurin-9-yl) propionylamino] benzoic acid ethyl ester.
68. The method of claim 3 wherein the compound is 4-[2-(6-oxo-2-phenyl-1,6-dihydropurin-9-yl) propionylamino] benzoic acid ethyl ester.
69. The method of claim 3 wherein the compound is 4-[[3-(6-chloropurin-9-yl) propionyl] methylamino] benzoic acid methyl ester.
70. The method of claim 3 wherein the compound is 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-[3-(2-oxopyrrolidin-1-yl)propyl] propanamide.
71. The method of claim 3 wherein the compound is 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-{2-[2-(2-oxopyrrolidin-1-yl)acetylamin]ethyl} propanamide.

72. The method of claim 3 wherein the compound is N-[3-(2-oxopyrrolidin-1-yl)propyl]-3-(6-oxo-2-thioxo-1,2,3,6-tetrahydropurin-9-yl) propanamide.

73. The method of claim 3 wherein the compound is 3-(1-benzyl-6-oxo-1,6-dihydropurin-9-yl)-N-(3-morpholin-4-yl)propyl propionamide.

74. The method of claim 1 wherein the compound is a tetrahydroindolone derivative or analogue where A is a 9-atom bicyclic moiety in which the 5-membered ring has one to three nitrogen atoms, the bicyclic moiety of Formula (XIV)



where:

- (a) N_1 is bonded to L;
- (b) A_2 and A_3 are C or N;
 - (i) If A_2 and A_3 are both C and the bond between A_2 and A_3 is a single bond, then the bond between A_2 and R_2 is two single bonds, two hydrogen atoms or is a double bond in which R_2 is O or S and R_3 is two hydrogen atoms;
 - (ii) If A_2 and A_3 are both C and the bond between A_2 and A_3 is a double bond, then R_3 is hydrogen, the bond between A_2 and R_2 is a single bond and R_2 is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;
 - (iii) If A_2 and A_3 are both N, then the bond between A_2 and A_3 is a double bond and R_2 and R_3 are not present;
 - (iv) If A_2 is N and A_3 is C, then the bond between A_2 and A_3 is a double bond, R_2 is not present, and R_3 is hydrogen;
 - (v) If A_2 is C, A_3 is N, and the bond between A_2 and A_3 is a double bond, then R_3 is not present, the bond between A_2 and R_2 is a single bond, and R_2 is hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, or heteroaralkenyl;

(vi) If A_2 is C, A_3 is N, and the bond between A_2 and A_3 is a single bond, then R_3 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, or heteroaralkenyl, the bond between A_2 and R_2 is a double bond, and A_2 is O or S;

(c) R_5 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NHQ_1 , NQ_1Q_2 , OH , OQ_1 , or SQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q_1 and Q_2 are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom, which can be N, O, or S, of which the N can be further substituted with Y_2 , where Y_2 is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(d) R_5 is hydrogen unless R_5 is alkyl, in which case R_5 is hydrogen or the same alkyl as R_5 ;

(e) R_5 and R_5 can be taken together as a double bond to C_5 , and can be O, S, NQ_3 , or C which can be substituted with one or two groups R_5 , where Q_3 is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(f) R_6 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, NH_2 , NHQ_4 , NQ_4Q_5 , OH , OQ_4 , or SQ_4 , where Q_4 and Q_5 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl,

alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₄ and Q₅ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom, which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;


(g) R₆ is hydrogen unless R₆ is alkyl, in which case R₆ is hydrogen or the same alkyl as R₆;

(h) R₆ and R₆ can be taken together as a double bond to C₆ and can be O, S, NQ₆, or C which can be substituted with one or two groups R₅, and where Q₆ is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and

(i) R₇ is hydrogen unless R₅ is alkyl and R₅ is hydrogen, in which case R₇ is the same alkyl as R₅.

75. The method of claim 74 wherein A is a tetrahydroindolone moiety.

76. The method of claim 75 wherein the tetrahydroindolone moiety is a tetrahydroindolone moiety of formula (XV)



in which:

(a) R_5 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NH_1 , NQ_1Q_2 , OH , OQ_1 , or SQ_1 , where Q_1 and Q_2 are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S;

(b) $R_{5'}$ is hydrogen;

(c) R_6 is hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, NH_2 , NHW_1 , NQ_1Q_2 , OH , OQ_1 , or SQ_1 , where Q_1 and Q_2 are aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S and where W_1 is alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from one to three heteroatoms which can be N, O, or S;

(d) R is hydrogen; and


(e) R_7 is hydrogen.

77. The method of claim 76 wherein R_5 , $R_{5'}$, R_6 , $R_{6'}$, and R_7 are all hydrogen.

78. The method of claim 77 wherein the compound is 4-[3-(4-oxo-4,5,6,7-tetrahydroindolon-1-yl) propionylamino] benzoic acid ethyl ester.

79. The method of claim 77 wherein the compound is 4-[3-(4-oxo-4,5,6,7-tetrahydroindolon-1-yl) propionylamino] benzoic acid.

80. The method of claim 1 wherein A is an amino-substituted 6-membered heterocyclic moiety of formula (XVI)



where:

(a) if the bond between N₁ and the bond between C₆ is a single bond, then the bond between C₆ and R₆ is a double bond, R₆ is O or S, and R₁ is hydrogen, alkyl, aralkyl, cycloalkyl, or heteroaralkyl;

(b) if the bond between N₁ and C₆ is a double bond, then the bond between C₆ and R₆ is a single bond, R₁ is not present, and R₆ is hydrogen, halo, amino, OH, OQ₁, SQ₁, NHNH₂, NQ₁Q₂, or NHQ₁, where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(c) if the bond between C₂ and N₃ is a single bond, then the bond between C₂ and R₂ is a double bond, R₂ is O or S, and R₃ is hydrogen or alkyl;

(d) if the bond between C₂ and N₃ is a double bond, then the bond between C₂ and R₂ is a single bond, R₃ is not present, and R₂ is hydrogen, alkyl, aralkyl, cycloalkyl, heteroaralkyl, halo, amino, OH, OQ₁, SQ₁, NHNH₂, NHOQ₁, NQ₁Q₂, or NHQ₁, where Q₁ and Q₂ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl,

aroyl, aralkanoyl, heteroaralkanoyl, heteroaroyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, or heteroaralkylsulfonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₁ and Q₂ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₃, where Y₃ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S;

(e) R₄ is hydrogen, alkyl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, or heteroarylaminocarbonyl;

(f) A₅ is carbon or nitrogen;

(g) if A₅ is nitrogen, then R₅ is not present;

(h) if A₅ is carbon, then R₅ is hydrogen, amino, alkyl, alkoxy, halo, nitro, aryl, cyano, alkenyl, or alkaryl;

(i) if R₅ and R₆ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S; and

(j) N₄ is bonded to L.

81. The method of claim 80 wherein A_5 is carbon and the 6-membered heterocyclic moiety is a pyrimidine moiety.
82. The method of claim 81 wherein R_2 is O and R_3 is hydrogen.
83. The method of claim 82 wherein the pyrimidine moiety is selected from the group consisting of cytosine, thymine, uracil, 3-methyluracil, 3-methylthymine, 4-methylcytosine, 5-methylcytosine, 5-hydroxymethylcytosine, 5-hydroxyuracil, 5-carboxymethyluracil, and 5-hydroxymethyluracil.
84. The method of claim 81 wherein R_2 is S and R_3 is hydrogen.
85. The method of claim 84 wherein the pyrimidine moiety is selected from the group consisting of 2-thiouracil, 5-methylamino-2-thiouracil, 5-methyl-2-thiouracil, 2-thiocytosine.
86. The method of claim 81 wherein R_2 is amino and the bond between C_2 and N_3 is a double bond.
87. The method of claim 86 wherein the pyrimidine moiety is selected from the group consisting of 2-aminopyrimidinone and 2-amino-4-chloropyrimidine.
88. The method of claim 81 wherein R_2 is hydrogen and the bond between C_2 and N_3 is a double bond.
89. The method of claim 88 wherein the pyrimidine moiety is selected from the group consisting of 4-chloropyrimidine, 5-amino-4-chloropyrimidine, 4-chloro-5-methylpyrimidine, 4-chloro-5-hydroxymethylpyrimidine, and 4-chloro-5-carboxymethylpyrimidine.
90. The method of claim 81 wherein R_1 is hydrogen, methyl, or ethyl, R_5 is hydrogen, methyl, or ethyl, and R_6 is O.
91. The method of claim 90 wherein the pyrimidine moiety is pyrimidinone.
92. The method of claim 81 wherein the compound is 4-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.
93. The method of claim 81 wherein the compound is 4-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.
94. The method of claim 81 wherein the compound is 4-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.

95. The method of claim 81 wherein the compound is 4-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

96. The method of claim 81 wherein the compound is 4-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

97. The method of claim 81 wherein the compound is 4-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

98. The method of claim 81 wherein the compound is 3-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.

99. The method of claim 81 wherein the compound is 3-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.

100. The method of claim 81 wherein the compound is 3-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid ethyl ester.

101. The method of claim 81 wherein the compound is 3-[3-(2-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

102. The method of claim 81 wherein the compound is 3-[3-(6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

103. The method of claim 81 wherein the compound is 3-[3-(5-amino-6-chloropyrimidin-4-ylamino) propionylamino] benzoic acid.

104. The method of claim 1 wherein L has the structure $-(CH_2)_n-$ wherein n is an integer from 1 to 6.

105. The method of claim 104 wherein n is 2.

106. The method of claim 104 wherein n is 3.

107. The method of claim 1 wherein the moiety B is $-OZ$.

108. The method of claim 107 wherein Z is hydrogen.

109. The method of claim 107 wherein Z is alkyl.

110. The method of claim 109 wherein Z is selected from the group consisting of methyl, ethyl, butyl, propyl, and isopropyl.

111. The method of claim 1 wherein B is $-N(Y_1)-D$.


112. The method of claim 111 wherein Y_1 is hydrogen.

113. The method of claim 111 wherein Y_1 is lower alkyl.

114. The method of claim 113 wherein Y_1 is methyl.

115. The method of claim 111 wherein D is a moiety having at least one polar, charged, or hydrogen-bond-forming group to increase the water-solubility of the compound.

116. The method of claim 115 wherein D is a carboxylic acid or carboxylic acid ester with the structure




wherein p is an integer from 1 to 6 and W_1 is selected from the group consisting of hydrogen and lower alkyl.

117. The method of claim 116 wherein W_1 is hydrogen.


118. The method of claim 116 wherein W_1 is ethyl.

119. The method of claim 115 wherein D and Y_1 are taken together to form a piperazine derivative of the structure



wherein Q_1 is hydrogen, methyl, ethyl, butyl, or propyl, and Q_2 is hydrogen or methyl, where, if Q_2 is methyl, it can be located on either of the two possible positions in the piperazine ring.


120. The method of claim 115 wherein D has the structure



wherein one of Z_1 and Z_2 is hydrogen and the other is Z_1 and Z_2 is ---COOH or ---COOW_1 , wherein W_1 is alkyl.


121. The method of claim 120 wherein W_1 is selected from the group consisting of methyl, ethyl, propyl, butyl, and isobutyl.

122. The method of claim 115 wherein D is a phenylsulfonamidyl moiety of the structure




wherein p is an integer from 0 to 6.

123. The method of claim 115 wherein D is an alkylpyridyl moiety of the structure



wherein p is an integer from 1 to 6.

124. The method of claim 114 wherein D is an dialkylaminoalkyl moiety of the structure



wherein p is an integer from 1 to 6 and Q₇ and Q₈ are alkyl, aralkyl, heteroaralkyl, aryl, heteroaryl, alkanoyl, aroyl, aralkanoyl, heteroaralkanoyl, or heteroaroyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S, and when Q₇ and Q₈ are present together and are alkyl, they can be taken together to form a 5- or 6-membered ring which can contain one other heteroatom which can be N, O, or S, of which the N can be further substituted with Y₂, where Y₂ is alkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, alkanoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, aralkylaminocarbonyl, or heteroaralkylaminocarbonyl, in which the alkyl portions can be cyclic and can contain from 1 to 3 heteroatoms which can be N, O, or S.

125. The method of claim 124 wherein Q₇ and Q₈ are each alkyl.

126. The method of claim 125 wherein Q₇ and Q₈ are each selected from the group consisting of methyl, ethyl, propyl, butyl, and isobutyl.


127. The method of claim 126 wherein Q₇ and Q₈ are taken together to form 5- or 6-membered optionally substituted ring.

128. The method of claim 127 wherein the ring is a morpholinyl ring.

129. The method of claim 127 wherein the ring is a pyrrolidinyl ring that is optionally substituted with oxo.

130. The method of claim 126 wherein the ring is a piperidinyl ring that is optionally substituted with methyl or ethyl.

131. The method of claim 115 wherein D is an alkylpyrrolidinyl moiety of the structure



wherein p is an integer from 1 to 6 and W_1 is selected from the group consisting of methyl, ethyl, and propyl.

132. The method of claim 1 wherein the compound has a log P of from about 1 to about 4.

133. The method of claim 1 wherein the action of the compound having activity against drug-induced neuropathy is to induce upregulation of neurotrophic factor synthesis.

134. The method of claim 133 wherein the neurotrophic factor is selected from the group consisting of NGF, NT-3, BDNF, and NT-4/5.

135. The method of claim 1 wherein the administration of the compound having activity against drug-induced neuropathy induces peripheral nerve sprouting in the skin of the patient to whom the compound was administered.

136. The method of claim 135 wherein the peripheral nerve sprouting is nociceptive nerve sprouting.

137. The method of claim 136 wherein the nociceptive nerve sprouting is induced without the occurrence of hyperalgesia.

138. The method of claim 1 wherein the drug-induced neuropathy is induced by the administration of an oncolytic drug.

139. The method of claim 138 wherein the oncolytic drug is selected from the group consisting of a vinca alkaloid, cisplatin, paclitaxel, suramin, altretamine,

carboplatin, chlorambucil, cytarabine, dacarbazine, docetaxel, etoposide, fludarabine, ifosfamide with mesna, tamoxifen, teniposide, and thioguanine.

140. The method of claim 139 wherein the oncolytic drug is vincristine.

141. The method of claim 139 wherein the oncolytic drug is paclitaxel.

142. The method of claim 139 wherein the oncolytic drug is cisplatin.

1/3




FIG. 1




FIG. 2

2/3

Sensory Nerve Conduction Velocity Effects of AIT-082 on Vincristine-Induced Neuropathy




FIG. 3

EMG Amplitude Effects of AIT-082 on Vincristine-Induced Neuropathy




FIG. 4

3/3




FIG. 5